

# Spotted Morning Thrush

*Cichladusa guttata*

616 plots from 1,894 records

Virgin Yellow-vented Bulbul squares **15**

Single YvB month **32**

Two YvB months **34**

### Seasonality


	presence	bs	eggs
January	171	7	3
February	171	4	5
March	142	4	3
April	131	16	2
May	136	5	3
June	136	5	1
July	155	1	-
August	163	3	-
September	137	2	1
October	161	7	4
November	132	5	4
December	165	8	3

### Species

- Presence
- Breeding season
- Month with eggs

### Elevation (m)

- 1 - 300
- 300 - 500
- 500 - 700
- 700 - 900
- 900 - 1,100
- 1,100 - 1,300
- 1,300 - 1,500
- 1,500 - 1,700
- 1,700 - 1,900
- 1,900 - 2,100



with support from  
THE LEVERHULME TRUST

**Tanzania Bird Atlas  
Preliminary Map  
October 2006  
From 735,041  
Database records**

J F M A  
M J J A  
S O N D  
# # # #

Monthly records  
within each square

Britton (1980) gives 9 localities for the nominate race shown here as red circles and 5 localities for the race *C. g. rufipennis* shown here as blue squares.

There appears to be an expansion of range northwards and westwards from the Wembere to the eastern shores of Lake Victoria and well into Sukumaland.

An indication perhaps of habitat degradation and a gradual drying of the climate.

29° 30° 31° 32° 33° 34° 35° 36° 37° 38° 39° 40°